Table 11. Estimated numbers of persons living with AIDS at the end of 2004, by race/ethnicity, sex, and transmission category—United States

Transmission category	White, not Hispanic		Black, not Hispanic		Hispanic		Asian/Pacific Islander		American Indian/Alaska Native		Totala	
	No.	<u>*************************************</u>	No.	*************************************	No.	/////////////////////////////////////	No.	<u>"""</u>	No.	<u>"""</u>	No.	%
Male adult or adolescent												
Male-to-male sexual contact	95,737	75	52,120	44	33,717	52	2,445	72	645	56	185,326	58
Injection drug use	13,137	10	34,797	29	18,472	28	314	9	190	17	67,091	21
Male-to-male sexual contact and	,		- 1,1 - 1		,			_			,	
injection drug use	11,366	9	9,174	8	4,361	7	162	5	195	17	25,367	8
Heterosexual contact	5,291	4	21,565	18	8,204	13	387	11	96	8	35,671	11
Other ^b	1,854	1	1,654	1	619	1	76	2	17	1	4,242	1
Subtotal	127,385	100	119,310	100	65,373	100	3,384	100	1,141	100	317,698	100
Female adult or adolescent												
Injection drug use	7,147	40	18,184	32	5,802	33	102	16	136	39	31,472	34
Heterosexual contact	10,313	57	36,791	65	11,561	65	491	76	203	58	59,599	64
Other ^b	529	3	1,474	3	416	2	51	8	11	3	2,494	3
Subtotal	17,990	100	56,449	100	17,780	100	644	100	350	100	93,566	100
Child (<13 yrs at diagnosis)												
Perinatal	508	91	2,435	98	809	95	16	93	14	99	3,797	97
Other ^c	51	9	39	2	39	5	1	7	0	1	131	3
Subtotal	559	100	2,474	100	848	100	17	100	14	100	3,927	100
Totald	145,935		178,233		84,001		4,045		1,506		415,193	

Note. These numbers do not represent reported case counts. Rather, these numbers are point estimates, which result from adjustments of reported case counts. The reported case counts are adjusted for reporting delays and for redistribution of cases in persons initially reported without an identified risk factor. The estimates do not include adjustment for incomplete reporting.

^a Includes 1,472 persons of unknown race or multiple races.

^b Includes hemophilia, blood transfusion, perinatal, and risk factor not reported or not identified.

^c Includes hemophilia, blood transfusion, and risk factor not reported or not identified.

^d Includes 2 person of unknown sex. Because column totals were calculated independently of the values for the subpopulations, the values in each column may not sum to the column total.